

# UNCLASSIFIED

AD NUMBER
AD834065
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; 08 AUG 1967. Other requests shall be referred to FForeign Technology Division, Attn: TDBDP, Wright-Patterson AFB, OH 45433-6508.
AUTHORITY
FTD, USAF ltr, 7 Oct 1971

THIS PAGE IS UNCLASSIFIED

AD834065

FTD-HT-67-73

## FOREIGN TECHNOLOGY DIVISION



ON THE INFLUENCE OF REPEATED NOISE STRESS ON RATS


by

V. Hrubec and V. Benes


CLASSIFICATION: UNCLASSIFIED

This document is subject to special export controls and is not to be released to foreign governments or foreign nationals without prior approval of the Foreign Technology Division.

EDWARD P. W. R. APPROVED 45433



FOREIGN TECHNOLOGY DIVISION



Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public.



## UNEDITED ROUGH DRAFT TRANSLATION

ON THE INFLUENCE OF REPEATED NOISE STRESS  
ON RATS

By: V. Hrubyš and V. Beneš

English pages: 6

SOURCE: Acta Biologica et Medica Germanica. (German  
Transactions in Biology and Medicine), Vol. 15,  
No. 5, 1965, pp. 592-596.

Translated by: E. Novak/TDBXT

GE/0020-65-015-005

TP7001007

THIS TRANSLATION IS A RENDERING OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:  
TRANSLATION DIVISION  
FOREIGN TECHNOLOGY DIVISION  
WP-APB, ONR.

THE RECORD

NEW SERIES

2

01 Acc Nr TP7001007		68 Translation Nr HT6700073		65 X Ref Acc Nr		76 Reel/Frame Nr 1881 0308	
97 Header Clas U..CL		63 Clas UNCL, 0		64 Control Markings 0		94 Expansion 40 Ctry Info GE	
02 Ctry GE	03 Ref 0020	04 Yr 65	05 Vol 015	06 Iss 005	07 B. Pg. 0592	45 E. Pg. 0596	10 Date NONE

Transliterated Title

UBER DEN EINFLUSS WIEDERHOLTER LARMBELASTUNG AUF RATTEN

09 English Title

ON THE INFLUENCE OF REPEATED NOISE STRESS ON RATS

43 Source

ACTA BIOLOGICA ET MEDICA GERMANICA (GERMAN)

42 Author

HRUBES, V.

16 Co-Author

BENES, V.

16 Co-Author

NONE

16 Co-Author

NONE

16 Co-Author

NONE

98 Document Location

47 Subject Codes

06

39 Topic Tags:

fatty acid, adrenal gland, blood plasma, catecholamine

**ABSTRACT:** The influence of the effect of repeated noise on rats was investigated. In comparison with untreated control animals and the initial values of the test animals, it was determined that the excretion of catechol amines in the urine is increased, free fatty-acids in the blood plasma increases, there is an increase in the weight of the adrenal glands and the growth of the animal is inhibited. The development of the weight curves also displays characteristic changes. The secretion of 5-hydroxyindolic acetic acid, however, does not change significantly. Consequently, noise is to be used as a model for the study of stress reactions. English Translation: 6 pages.

**BLANK PAGE**

## ON THE INFLUENCE OF REPEATED NOISE STRESS ON RATS

V. Hruběš and V. Beneš

### Summary

The influence of the effect of repeated noise on rats was investigated. In comparison to untreated control animals and the initial values of the test animals, it was determined that the excretion of catechol amines in the urine is increased, free fatty-acids in the blood plasma increases, there is an increase in the weight of the adrenal glands and the growth of the animal is inhibited. The development of the weight curves also displays characteristic changes. The secretion of 5-hydroxyindolic acetic acid, however, does not change significantly.

Noise not only effects the hearing, but also the central nervous system and, in so doing, has an influence on subordinate systems which results in corresponding stress reactions.

The reaction of the organism to a stress is a result of a simultaneous activation of the sympathoadrenales and the humoral systems, especially with regard to changes in the action of the adrenal cortex. In comparison, catechol amine is at times active as a neuro-chemical mediator in various regulative functions, and also ACTH which is probably liberated under the influence of the stimulation from the hypothalamus (1, 2). Due to the influence of the catechol amine, glucose is mobilized from glycogen and free fatty-acids from the fat deposits. Nevertheless, ACTH takes

place in the metabolism of carbohydrates and fats. Hyperglycaemia is the primary result which is relieved by a hyperlipaemia in another phase. Catechol amine and its satellites are secreted in ever-increasing amounts in the urine.

In our article we concentrate on a number of important components that always take place in the stress reaction, namely, the catechol amine, serotonin and the free fatty-acids in the plasma.

### METHODS

Female rats having an initial weight of from 132 to 150 grams that were fed a standard diet were used as test animals. The rats were subjected to the influence of noise for 6 hours a day for 4 days (noise was amplified by means of a transistor generator having an intensity of 95 db); this was divided by a 3 day rest period. Two test series were made up with 10 experimental animals and 10 control animals. The animals in the first series (A) were tested for 2 weeks and the second group (B), 4 weeks. In all animals the body weight was regularly checked. The amount of catechol amine was determined constantly by the spectral photometric method according to Oesterling and Tse (3) and 5-hydroxy indolic acid (5-HIES) was determined colorimetrically according to Udenfriend and associates (4). At the end of the test the animals were killed, the adrenals were recovered and the fatty acids that were not oesterified in the plasma were determined according to Dole (5).

### RESULTS AND DISCUSSION

The response time of the catechol amine excretion is shown in Fig. 1 expressed in extinction values. From this we can see that in the four week experiment, there is a clear increase in the excretion, namely, in comparison to the initial values as well as also to the values in the control group. On the other hand, when looking



for 5 HIES there was no essential difference between the test and control animals. The amount of the free fatty-acids in the plasma is shown in Fig. 2. In both cases higher values were obtained in the test rats than in the control animals. The weight curves are shown in Fig. 3. From the illustration, we can see that the exposed rats gained weight slower. When the exposure time and the resting time was exchanged during a single week, it was especially determined in test B that during the exposure time the rats always suffered from a statistical significant retardation of their growth and in the last period of time a significant weight loss. Partial comparisons of the increase in weight under both time conditions (noise, quiet) during the same week statistically significant differences are shown (Table 1).




Fig. 1. Transient response of catecholamine excretion (in extinction values).  
The long solid line = experiment; the weaker line = control.

In comparing the growth curve of control and test rats, we can see that in the last three weeks the difference in the weight increase during the exposure is statistically significant ( $P < 0.01$ ).

The weight of the adrenals, that is,

the absolute weight as well as the weight based on the 100 g rat weight is shown in Fig. 4. The adrenals of the test rats were, on the average, heavier; however,




Fig. 2. Free fatty-acid content in plasma.

the differences are not significant from a statistic standpoint.




Abb. 3. Gewicht der Versuchstiere.

— = Experiment; - - - = Kontrolle

Fig. 3. Weight of the test animals.

The solid line = experimental; the weaker line = control.

**KEY:** (1) absolute increase; (2) transient response; (3) average weight; (4) noise.

Table 1.

Increase in Weight in the Test and Control of Rats, Experiment A.

(1) Lärm		(2) Kontrolle		t
$\bar{x}$	$t s_{\bar{x}}$	$\bar{x}$	$t s_{\bar{x}}$	
(3) d <sub>1</sub> Ruhe 2,500 ± 2,65		d <sub>1</sub> Ruhe 2,875 ± 1,59		0,27
(4) d <sub>2</sub> Lärm 0,0 ± 2,65		d <sub>2</sub> Ruhe -1,085 ± 1,59		0,79

Experiment B

(1) Lärm		(2) Kontrolle		t
$\bar{x}$	$t s_{\bar{x}}$	$\bar{x}$	$t s_{\bar{x}}$	
(3) d <sub>1</sub> Ruhe 1,625 ± 1,43		d <sub>1</sub> Ruhe 2,250 ± 0,95		0,82
(4) d <sub>2</sub> Lärm 1,750 ± 1,43		d <sub>2</sub> Ruhe -0,459 ± 0,95		2,90
d <sub>3</sub> Lärm 0,223 ± 1,43		d <sub>3</sub> Ruhe -0,376 ± 0,95		0,78
d <sub>4</sub> Lärm 1,375 ± 1,43		d <sub>4</sub> Ruhe -0,250 ± 0,95		2,13

Experiment B

(5) Zusammenfassung von d<sub>2</sub> + d<sub>3</sub> + d<sub>4</sub> (nur während der Expositionszeit)

Lärm		Kontrolle		t
$\bar{x}$	$t s_{\bar{x}}$	$\bar{x}$	$t s_{\bar{x}}$	
d <sub>2</sub> + d <sub>3</sub> + d <sub>4</sub> 1,12 ± 0,6980		d <sub>2</sub> + d <sub>3</sub> + d <sub>4</sub> -0,36 ± 0,49		3,55

**KEY:** (1) noise; (2) control; (3) quiet; (4) noise; (5) combination of d<sub>2</sub> + d<sub>3</sub> + d<sub>4</sub> (only during the exposure time).




Abb. 4. Nebennieren-Gewicht.  
K = Kontrolle; E = Versuchstiere

Fig. 4. The weight of the adrenals.  
**KEY:** (1) absolute average weight;  
(2) average weight. K = control; E =  
the test animals.

The results of the biochemical investigations are in agreement with those of the other authors. The results on the biological changes proves the work that was done in America on the experimental audiogenic stress (6), (7). From the present work and the results that we obtained earlier (8) - (10), it can be stated that, due to the influence of noise, similar metabolic reactions can occur as those in tests in which other stress types were used. Consequently, noise is to be used as a model for the study of stress reactions throughout.

Address of the authors: V. Hruběš and V. Beneš, 48 Štrobářova, Prague 10.

### Literature

- [1] CHARVÁT, J.: Čas. lék. česk. 103, 761 (1964).
- [2] SCHREIBER, V.: Čas. lék. česk. 103, 766 (1964).
- [3] OESTERLING, M. J. u. R. L. TSE: Amer. J. med. Technol. 27, 112 (1961).
- [4] UDENFRIEND, S., E. TITUS u. H. WEISSBACH: J. biol. Chemistry 216, 499 (1955).
- [5] DOLE, V. P.: J. clin. Invest. 35, 150 (1956).
- [6] SACKLER, A. M., A. S. WELTMAN, M. BRADSHAW u. P. JR. JURTSUK: Acta endocrinol. (Copenhagen) 31, 405 (1959).